

Yrast level structure and decay of ^{137}I and neighboring nuclei from spontaneous fission of ^{252}Cf

J. Gilat, S.C. Wu†, J.O. Rasmussen, R.B. Firestone, S.J. Asztalos*

Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

M.W. Simon, C.Y. Wu, D. Cline

Nuclear Structure Research Laboratory, University of Rochester, Rochester, NY 14623, USA

M.A. Stoyer, Y.X. Dardenne

Lawrence Livermore National Laboratory, Livermore, CA 94550, USA

J.H. Hamilton, A.V. Ramayya

Physics Department, Vanderbilt University, Nashville, TN 37235, USA

We have continued the spectroscopic analysis of coincident prompt γ -ray data collected in a thick ^{252}Cf source Gammasphere experiment carried out in December 1995, focusing on odd Z nuclei. The results were combined with those obtained with a thin ^{252}Cf source, measurement of γ -rays emitted in flight and simultaneous detection of both fission fragments using the CHICO PPAC array inside Gammasphere [1]. The data from the two experiments are complementary. The identification of coincident fission-partner γ -rays in the thick source spectra facilitates the initial isotope assignment of observed γ -ray cascades. Furthermore, the energy resolution of these spectra is considerably better than that of the Doppler-corrected thin source data. On the other hand, the mass resolution inherent to the thin source experiment results in lower background from partner γ -rays and from randomly overlapping transitions in other fission products. This in turn leads to less ambiguous identification and assignment of weaker γ -rays and enables the extension of level schemes to higher spin values. Moreover, the ~ 10 ns flight time can be used to estimate level lifetimes and γ -ray multipolarities.

We used data from the two experiments to identify and assign γ -transitions in $^{134,136}\text{Te}$ and in $^{135-138}\text{I}$, and to elucidate yrast level schemes for these isotopes, few nucleons above the doubly magic ^{132}Sn . In ^{137}I we identified 16 new γ -transitions between 186- and 862 keV and placed them in a level scheme with two (positive parity?) bands, extending up to spin of $37/2$. There is no

overlap between these yrast levels and levels previously observed in β -decay of ^{137}Te [2]. The ground-state band of ^{137}I is very similar to that reported for the even-even ^{136}Te [3], and seems to reflect a relatively loose coupling of a $g_{7/2}$ proton to the Te core. We have also been able to extend the ^{136}Te level scheme of Ref. [3] up to $J=16$, and to identify a number of side transitions in this nucleus. The level scheme we constructed for ^{136}I , confirms and extends the recently published [4] scheme, based on a Eurogam study of ^{248}Cm spontaneous fission. The similarity of the main features of this level scheme to those of ^{135}Te again indicates the loose coupling of the 53^{rd} proton to Te cores. The energies of the uppermost γ -emitting levels in ^{136}I and ^{135}Te exceed the neutron binding energies of these nuclei, reflecting the angular momentum hindrance of neutron emission near yrast.

Footnotes and References

* On sabbatical leave from Soreq Nuclear Research Center, Yavne 81800, Israel.

† On sabbatical leave from National Tsing Hua University, Hsinchu, Taiwan 30043.

1. M.W. Simon et al., Proceedings of the International Conference on Fission and Properties of Neutron-Rich Nuclei, held Nov. 10-15, 1997 on Sanibel Island, FL (in print)
2. M. Samri et al., *Z. Phys.* **A321** (1985) 255
3. K. Butler-Moore et al., *J. Phys. (London)* **G19** (1993) L121
4. P. Bhattacharyya et al., *Phys. Rev.* **C56** (1997) R2363